

IOWA STATE UNIVERSITY

Digital Repository

Physics and Astronomy Publications

Physics and Astronomy

4-2010

Enhanced Production of Direct Photons in Au plus Au Collisions at $\sqrt{s(NN)}=200$ GeV and Implications for the Initial Temperature

Andrew Adare

University of Colorado, Boulder

Sergey Belikov

Iowa State University

Paul Constantin

Iowa State University

Nathan C. Grau

Iowa State University

John C. Hill

Iowa State University, jhill@iastate.edu

See next page for additional authors

Follow this and additional works at: http://lib.dr.iastate.edu/physastro_pubs



Part of the [Elementary Particles and Fields and String Theory Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/physastro_pubs/313. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Physics and Astronomy at Iowa State University Digital Repository. It has been accepted for inclusion in Physics and Astronomy Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Enhanced Production of Direct Photons in Au plus Au Collisions at $\sqrt{s(NN)}=200$ GeV and Implications for the Initial Temperature

Abstract

The production of $e^{+}e^{-}$ pairs for $m(e^{+}e^{-}) < 0.3$ GeV/c(2) and $1 < p(T) < 5$ GeV/c is measured in p + p and Au + Au collisions at $\sqrt{s(NN)} = 200$ GeV. An enhanced yield above hadronic sources is observed. Treating the excess as photon internal conversions, the invariant yield of direct photons is deduced. In central Au + Au collisions, the excess of the direct photon yield over p + p is exponential in transverse momentum, with an inverse slope $T = 221 \pm 19(\text{stat}) \pm 19(\text{syst})$ MeV. Hydrodynamical models with initial temperatures ranging from T_{init} similar to 300-600 MeV at times of similar to 0.6-0.15 fm/c after the collision are in qualitative agreement with the data. Lattice QCD predicts a phase transition to quark gluon plasma at similar to 170 MeV.

Disciplines

Elementary Particles and Fields and String Theory | Physics

Comments

This article is published as Adare, Ad, S. Afanasiev, C. Aidala, N. N. Ajitanand, Y. Akiba, H. Al-Bataineh, J. Alexander et al. "Enhanced production of direct photons in Au+ Au collisions at $\sqrt{s(NN)} = 200$ GeV and implications for the initial temperature." *Physical Review Letters* 104, no. 13 (2010): 132301. DOI:[10.1103/PhysRevLett.104.132301](https://doi.org/10.1103/PhysRevLett.104.132301). Posted with permission.

Authors

Andrew Adare, Sergey Belikov, Paul Constantin, Nathan C. Grau, John C. Hill, John G. Lajoie, Alexandre Lebedev, Craig Ogilvie, H. Pei, Jan Rak, Marzia Rosati, S. Skutnik, Carla Vale, et al., and PHENIX Collaboration

Enhanced Production of Direct Photons in Au + Au Collisions at $\sqrt{s_{NN}} = 200$ GeV and Implications for the Initial Temperature

A. Adare,⁸ S. Afanasiev,²² C. Aidala,⁹ N. N. Ajitanand,⁴⁹ Y. Akiba,^{43,44} H. Al-Bataineh,³⁸ J. Alexander,⁴⁹ A. Al-Jamel,³⁸ K. Aoki,^{28,43} L. Aphecetche,⁵¹ R. Armendariz,³⁸ S. H. Aronson,³ J. Asai,⁴⁴ E. T. Atomssa,²⁹ R. Averbeck,⁵⁰ T. C. Awes,³⁹ B. Azmoun,³ V. Babintsev,¹⁸ G. Baksay,¹⁴ L. Baksay,¹⁴ A. Baldisseri,¹¹ K. N. Barish,⁴ P. D. Barnes,³¹ B. Bassalleck,³⁷ S. Bathe,⁴ S. Batsouli,^{9,39} V. Baublis,⁴² F. Bauer,⁴ A. Bazilevsky,³ S. Belikov,^{3,21,*} R. Bennett,⁵⁰ Y. Berdnikov,⁴⁶ A. A. Bickley,⁸ M. T. Bjornedal,⁹ J. G. Boissevain,³¹ H. Borel,¹¹ K. Boyle,⁵⁰ M. L. Brooks,³¹ D. S. Brown,³⁸ D. Bucher,³⁴ H. Buesching,³ V. Bumazhnov,¹⁸ G. Bunce,^{3,44} J. M. Burward-Hoy,³¹ S. Butsyk,^{31,50} S. Campbell,⁵⁰ J.-S. Chai,²³ B. S. Chang,⁵⁸ J.-L. Charvet,¹¹ S. Chernichenko,¹⁸ J. Chiba,²⁴ C. Y. Chi,⁹ M. Chiu,^{9,19} I. J. Choi,⁵⁸ T. Chujo,⁵⁵ P. Chung,⁴⁹ A. Churnyn,¹⁸ V. Cianciolo,³⁹ C. R. Clevén,¹⁶ Y. Cobigo,¹¹ B. A. Cole,⁹ M. P. Comets,⁴⁰ P. Constantin,^{21,31} M. Csanád,¹³ T. Csörgő,²⁵ T. Dahms,⁵⁰ K. Das,¹⁵ G. David,³ M. B. Deaton,¹ K. Dehmelt,¹⁴ H. Delagrangé,⁵¹ A. Denisov,¹⁸ D. d'Enterria,⁹ A. Deshpande,^{44,50} E. J. Desmond,³ O. Dietzsch,⁴⁷ A. Dion,⁵⁰ M. Donadelli,⁴⁷ J. L. Drachenberg,¹ O. Drapier,²⁹ A. Drees,⁵⁰ A. K. Dubey,⁵⁷ A. Durum,¹⁸ V. Dzhordzhadze,^{4,52} Y. V. Efremenko,³⁹ J. Egdemir,⁵⁰ F. Ellinghaus,⁸ W. S. Emam,⁴ A. Enokizono,^{17,30} H. En'yo,^{43,44} B. Espagnon,⁴⁰ S. Esumi,⁵⁴ K. O. Eyser,⁴ D. E. Fields,^{37,44} M. Finger,^{5,22} M. Finger, Jr.,^{5,22} F. Fleuret,²⁹ S. L. Fokin,²⁷ B. Forestier,³² Z. Fraenkel,^{57,*} J. E. Frantz,^{9,50} A. Franz,³ A. D. Frawley,¹⁵ K. Fujiwara,⁴³ Y. Fukao,^{28,43} S.-Y. Fung,⁴ T. Fusayasu,³⁶ S. Gadrat,³² I. Garishvili,⁵² F. Gastineau,⁵¹ M. Germain,⁵¹ A. Glenn,^{8,52} H. Gong,⁵⁰ M. Gonin,²⁹ J. Gosset,¹¹ Y. Goto,^{43,44} R. Granier de Cassagnac,²⁹ N. Grau,²¹ S. V. Greene,⁵⁵ M. Grosse Perdekamp,^{19,44} T. Gunji,⁷ H.-Å. Gustafsson,³³ T. Hachiya,^{17,43} A. Hadj Henni,⁵¹ C. Haegemann,³⁷ J. S. Haggerty,³ M. N. Hagiwara,¹ H. Hamagaki,⁷ R. Han,⁴¹ H. Harada,¹⁷ E. P. Hartouni,³⁰ K. Haruna,¹⁷ M. Harvey,³ E. Haslum,³³ K. Hasuko,⁴³ R. Hayano,⁷ M. Heffner,³⁰ T. K. Hemmick,⁵⁰ T. Hester,⁴ J. M. Heuser,⁴³ X. He,¹⁶ H. Hiejima,¹⁹ J. C. Hill,²¹ R. Hobbs,³⁷ M. Hohlmann,¹⁴ M. Holmes,⁵⁵ W. Holzmann,⁴⁹ K. Homma,¹⁷ B. Hong,²⁶ T. Horaguchi,^{43,53} D. Hornback,⁵² M. G. Hur,²³ T. Ichihara,^{43,44} K. Imai,^{28,43} M. Inaba,⁵⁴ Y. Inoue,^{45,43} D. Isenhowe,¹ L. Isenhowe,¹ M. Ishihara,⁴³ T. Isobe,⁷ M. Issah,⁴⁹ A. Isupov,²² B. V. Jacak,^{50,†} J. Jia,⁹ J. Jin,⁹ O. Jinnouchi,⁴⁴ B. M. Johnson,³ K. S. Joo,³⁵ D. Jouan,⁴⁰ F. Kajihara,^{7,43} S. Kametani,^{7,56} N. Kamihara,^{43,53} J. Kamin,⁵⁰ M. Kaneta,⁴⁴ J. H. Kang,⁵⁸ H. Kanou,^{43,53} T. Kawagishi,⁵⁴ D. Kawall,⁴⁴ A. V. Kazantsev,²⁷ S. Kelly,⁸ A. Khanzadeev,⁴² J. Kikuchi,⁵⁶ D. H. Kim,³⁵ D. J. Kim,⁵⁸ E. Kim,⁴⁸ Y.-S. Kim,²³ E. Kinney,⁸ A. Kiss,¹³ E. Kistenev,³ A. Kiyomichi,⁴³ J. Klay,³⁰ C. Klein-Boesing,³⁴ L. Kochenda,⁴² V. Kochetkov,¹⁸ B. Komkov,⁴² M. Konno,⁵⁴ D. Kotchetkov,⁴ A. Kozlov,⁵⁷ A. Král,¹⁰ A. Kravitz,⁹ P. J. Kroon,³ J. Kubart,^{5,20} G. J. Kunde,³¹ N. Kurihara,⁷ K. Kurita,^{45,43} M. J. Kweon,²⁶ Y. Kwon,^{52,58} G. S. Kyle,³⁸ R. Lacey,⁴⁹ Y.-S. Lai,⁹ J. G. Lajoie,²¹ A. Lebedev,²¹ Y. Le Bornec,⁴⁰ S. Leckey,⁵⁰ D. M. Lee,³¹ M. K. Lee,⁵⁸ T. Lee,⁴⁸ M. J. Leitch,³¹ M. A. L. Leite,⁴⁷ B. Lenzi,⁴⁷ H. Lim,⁴⁸ T. Liška,¹⁰ A. Litvinenko,²² M. X. Liu,³¹ X. Li,⁶ X. H. Li,⁴ B. Love,⁵⁵ D. Lynch,³ C. F. Maguire,⁵⁵ Y. I. Makdisi,³ A. Malakhov,²² M. D. Malik,³⁷ V. I. Manko,²⁷ Y. Mao,^{41,43} L. Mašek,^{5,20} H. Masui,⁵⁴ F. Matathias,^{9,50} M. C. McCain,¹⁹ M. McCumber,⁵⁰ P. L. McGaughey,³¹ Y. Miake,⁵⁴ P. Mikeš,^{5,20} K. Miki,⁵⁴ T. E. Miller,⁵⁵ A. Milov,⁵⁰ S. Mioduszewski,³ G. C. Mishra,¹⁶ M. Mishra,² J. T. Mitchell,³ M. Mitrovski,⁴⁹ A. Morreale,⁴ D. P. Morrison,³ J. M. Moss,³¹ T. V. Moukhanova,²⁷ D. Mukhopadhyay,⁵⁵ J. Murata,^{45,43} S. Nagamiya,²⁴ Y. Nagata,⁵⁴ J. L. Nagle,⁸ M. Naglis,⁵⁷ I. Nakagawa,^{43,44} Y. Nakamiya,¹⁷ T. Nakamura,¹⁷ K. Nakano,^{43,53} J. Newby,³⁰ M. Nguyen,⁵⁰ B. E. Norman,³¹ A. S. Nyanin,²⁷ J. Nystrand,³³ E. O'Brien,³ S. X. Oda,⁷ C. A. Ogilvie,²¹ H. Ohnishi,⁴³ I. D. Ojha,⁵⁵ H. Okada,^{28,43} K. Okada,⁴⁴ M. Oka,⁵⁴ O. O. Omiwade,¹ A. Oskarsson,³³ I. Otterlund,³³ M. Ouchida,¹⁷ K. Ozawa,⁷ R. Pak,³ D. Pal,⁵⁵ A. P. T. Palounek,³¹ V. Pantuev,⁵⁰ V. Papavassiliou,³⁸ J. Park,⁴⁸ W. J. Park,²⁶ S. F. Pate,³⁸ H. Pei,²¹ J.-C. Peng,¹⁹ H. Pereira,¹¹ V. Peresedov,²² D. Yu. Peressounko,²⁷ C. Pinkenburg,³ R. P. Pisani,³ M. L. Purschke,³ A. K. Purwar,^{31,50} H. Qu,¹⁶ J. Rak,^{21,37} A. Rakotozafindrabe,²⁹ I. Ravinovich,⁵⁷ K. F. Read,^{39,52} S. Rembeczki,¹⁴ M. Reuter,⁵⁰ K. Reygers,³⁴ V. Riabov,⁴² Y. Riabov,⁴² G. Roche,³² A. Romana,^{29,*} M. Rosati,²¹ S. S. E. Rosendahl,³³ P. Rosnet,³² P. Rukoyatkin,²² V. L. Rykov,⁴³ S. S. Ryu,⁵⁸ B. Sahlmueller,³⁴ N. Saito,^{28,43,44} T. Sakaguchi,^{3,7,56} S. Sakai,⁵⁴ H. Sakata,¹⁷ V. Samsonov,⁴² H. D. Sato,^{28,43} S. Sato,^{3,24,54} S. Sawada,²⁴ J. Seele,⁸ R. Seidl,¹⁹ V. Semenov,¹⁸ R. Seto,⁴ D. Sharma,⁵⁷ T. K. Shea,³ I. Shein,¹⁸ A. Shevel,^{42,49} T.-A. Shibata,^{43,53} K. Shigaki,¹⁷ M. Shimomura,⁵⁴ T. Shohjoh,⁵⁴ K. Shoji,^{28,43} A. Sickles,⁵⁰ C. L. Silva,⁴⁷ D. Silvermyr,³⁹ C. Silvestre,¹¹ K. S. Sim,²⁶ C. P. Singh,² V. Singh,²¹ S. Skutnik,²¹ M. Slunečka,^{5,22} W. C. Smith,¹ A. Soldatov,¹⁸ R. A. Soltz,³⁰ W. E. Sondheim,³¹ S. P. Sorensen,⁵² I. V. Sourikova,³ F. Staley,¹¹ P. W. Stankus,³⁹ E. Stenlund,³³ M. Stepanov,³⁸ A. Ster,²⁵ S. P. Stoll,³ T. Sugitate,¹⁷ C. Suire,⁴⁰ J. P. Sullivan,³¹ J. Sziklai,²⁵ T. Tabaru,⁴⁴ S. Takagi,⁵⁴ E. M. Takagui,⁴⁷ A. Taketani,^{43,44} K. H. Tanaka,²⁴ Y. Tanaka,³⁶ K. Tanida,^{43,44} M. J. Tannenbaum,³ A. Taranenko,⁴⁹ P. Tarján,¹² T. L. Thomas,³⁷

M. Togawa,^{28,43} A. Toia,⁵⁰ J. Tojo,⁴³ L. Tomášek,²⁰ H. Torii,⁴³ R. S. Towell,¹ V.-N. Tram,²⁹ I. Tserruya,⁵⁷
 Y. Tsuchimoto,^{17,43} S. K. Tuli,² H. Tydesjö,³³ N. Tyurin,¹⁸ C. Vale,²¹ H. Valle,⁵⁵ H. W. van Hecke,³¹ J. Velkovska,⁵⁵
 R. Vertesi,¹² A. A. Vinogradov,²⁷ M. Virius,¹⁰ V. Vrba,²⁰ E. Vznuzdaev,⁴² M. Wagner,^{28,43} D. Walker,⁵⁰ X. R. Wang,³⁸
 Y. Watanabe,^{43,44} J. Wessels,³⁴ S. N. White,³ N. Willis,⁴⁰ D. Winter,⁹ C. L. Woody,³ M. Wysocki,⁸ W. Xie,^{4,44}
 Y. L. Yamaguchi,⁵⁶ A. Yanovich,¹⁸ Z. Yasin,⁴ J. Ying,¹⁶ S. Yokkaichi,^{43,44} G. R. Young,³⁹ I. Younus,³⁷ I. E. Yushmanov,²⁷
 W. A. Zajc,⁹ O. Zaudtke,³⁴ C. Zhang,^{9,39} S. Zhou,⁶ J. Zimányi,^{25,*} and L. Zolin²²

(PHENIX Collaboration)

- ¹Abilene Christian University, Abilene, Texas 79699, USA
²Department of Physics, Banaras Hindu University, Varanasi 221005, India
³Brookhaven National Laboratory, Upton, New York 11973-5000, USA
⁴University of California-Riverside, Riverside, California 92521, USA
⁵Charles University, Ovocný trh 5, Praha 1, 116 36, Prague, Czech Republic
⁶China Institute of Atomic Energy (CIAE), Beijing, People's Republic of China
⁷Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan
⁸University of Colorado, Boulder, Colorado 80309, USA
⁹Columbia University, New York, New York 10027 and Nevis Laboratories, Irvington, New York 10533, USA
¹⁰Czech Technical University, Zikova 4, 166 36 Prague 6, Czech Republic
¹¹Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France
¹²Debrecen University, H-4010 Debrecen, Egyetem tér 1, Hungary
¹³ELTE, Eötvös Loránd University, H-1117 Budapest, Pázmány P. s. 1/A, Hungary
¹⁴Florida Institute of Technology, Melbourne, Florida 32901, USA
¹⁵Florida State University, Tallahassee, Florida 32306, USA
¹⁶Georgia State University, Atlanta, Georgia 30303, USA
¹⁷Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan
¹⁸IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia
¹⁹University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA
²⁰Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague 8, Czech Republic
²¹Iowa State University, Ames, Iowa 50011, USA
²²Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia
²³KAERI, Cyclotron Application Laboratory, Seoul, Korea
²⁴KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan
²⁵KFKI Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences (MTA KFKI RMKI),
 H-1525 Budapest 114, POBox 49, Budapest, Hungary
²⁶Korea University, Seoul, 136-701, Korea
²⁷Russian Research Center "Kurchatov Institute", Moscow, Russia
²⁸Kyoto University, Kyoto 606-8502, Japan
²⁹Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France
³⁰Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³¹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA
³²LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubiere Cedex, France
³³Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden
³⁴Institut für Kernphysik, University of Muenster, D-48149 Muenster, Germany
³⁵Myongji University, Yongin, Kyonggido 449-728, Korea
³⁶Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan
³⁷University of New Mexico, Albuquerque, New Mexico 87131, USA
³⁸New Mexico State University, Las Cruces, New Mexico 88003, USA
³⁹Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
⁴⁰IPN-Orsay, Université Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France
⁴¹Peking University, Beijing, People's Republic of China
⁴²PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia
⁴³RIKEN, The Institute of Physical and Chemical Research, Wako, Saitama 351-0198, Japan
⁴⁴RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973-5000, USA
⁴⁵Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan
⁴⁶Saint Petersburg State Polytechnic University, St. Petersburg, Russia
⁴⁷Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil
⁴⁸System Electronics Laboratory, Seoul National University, Seoul, Korea
⁴⁹Chemistry Department, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA

⁵⁰*Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York 11794, USA*⁵¹*SUBATECH (Ecole des Mines de Nantes, CNRS-IN2P3, Université de Nantes) BP 20722-44307, Nantes, France*⁵²*University of Tennessee, Knoxville, Tennessee 37996, USA*⁵³*Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan*⁵⁴*Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan*⁵⁵*Vanderbilt University, Nashville, Tennessee 37235, USA*⁵⁶*Waseda University, Advanced Research Institute for Science and Engineering, 17 Kikui-cho, Shinjuku-ku, Tokyo 162-0044, Japan*⁵⁷*Weizmann Institute, Rehovot 76100, Israel*⁵⁸*Yonsei University, IPAP, Seoul 120-749, Korea*

(Received 25 April 2008; revised manuscript received 2 December 2009; published 29 March 2010)

The production of e^+e^- pairs for $m_{e^+e^-} < 0.3$ GeV/ c^2 and $1 < p_T < 5$ GeV/ c is measured in $p + p$ and Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. An enhanced yield above hadronic sources is observed. Treating the excess as photon internal conversions, the invariant yield of direct photons is deduced. In central Au + Au collisions, the excess of the direct photon yield over $p + p$ is exponential in transverse momentum, with an inverse slope $T = 221 \pm 19^{\text{stat}} \pm 19^{\text{syst}}$ MeV. Hydrodynamical models with initial temperatures ranging from $T_{\text{init}} \sim 300\text{--}600$ MeV at times of $\sim 0.6\text{--}0.15$ fm/ c after the collision are in qualitative agreement with the data. Lattice QCD predicts a phase transition to quark gluon plasma at ~ 170 MeV.

DOI: 10.1103/PhysRevLett.104.132301

PACS numbers: 25.75.Cj, 12.38.Mh, 13.85.Qk, 21.65.Qr

Experimental results from the Relativistic Heavy Ion Collider (RHIC) have established the formation of dense partonic matter in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV [1]. The large energy loss of light quarks and gluons [2] as well as that of heavy quarks [3] indicates that the matter is very dense. The strong elliptic flow of light [4,5] and charmed [3] hadrons indicate rapid thermalization. Such a hot, dense medium should emit thermal radiation [6]; the partonic phase is predicted to be the dominant source of direct photons with $1 < p_T < 3$ GeV/ c in Au + Au collisions at RHIC [7].

The observation of thermal photons will allow the determination of the initial temperature of the matter. However, the measurement of direct photons for $1 < p_T < 3$ GeV/ c is notoriously difficult due to a large background from hadronic decay photons. Direct photons contribute only $\approx 10\%$ above the background photon yield [7]. In general, any source of high energy photons can also emit virtual photons, which convert to low-mass e^+e^- pairs. For example, gluon Compton scattering ($q + g \rightarrow q + \gamma$) has an associated process that produces low-mass e^+e^- pairs through internal conversion ($q + g \rightarrow q + \gamma^* \rightarrow q + e^+e^-$). Consequently, we search for “quasireal” virtual photons, which appear as low invariant mass e^+e^- pairs.

The relation between photon production and the associated e^+e^- pair production can be written as [8,9]

$$\frac{d^2 n_{ee}}{dm_{ee}} = \frac{2\alpha}{3\pi} \frac{1}{m_{ee}} \sqrt{1 - \frac{4m_e^2}{m_{ee}^2}} \left(1 + \frac{2m_e^2}{m_{ee}^2}\right) S dn_\gamma. \quad (1)$$

Here α is the fine structure constant, m_e and m_{ee} are the masses of the electron and the e^+e^- pair, respectively, and S is a process dependent factor that goes to 1 as $m_{ee} \rightarrow 0$ or $m_{ee} \ll p_T$. Equation (1) also describes the relation between the photons from hadron decays (e.g., π^0 , $\eta \rightarrow$

$\gamma\gamma$, and $\omega \rightarrow \gamma\pi^0$) and the e^+e^- pairs from Dalitz decays (π^0 , $\eta \rightarrow e^+e^- \gamma$ and $\omega \rightarrow e^+e^- \pi^0$). For π^0 and η , the factor S is given by $S = |F(m_{ee}^2)|^2 (1 - \frac{m_{ee}^2}{M_h^2})^3$ [10], where M_h is the meson mass and $F(m_{ee}^2)$ is the form factor.

The factor S for a hadron h is zero for $m_{ee} > M_h$. We exploit this cutoff to separate the direct photon signal from the hadronic background. Since 80% of the hadronic photons are from π^0 decays, the signal to background (S/B) ratio for the direct photon signal improves by a factor of 5 for $m_{ee} > M_{\pi^0} = 135$ MeV/ c^2 , thereby allowing a direct photon signal that is 10% of the background to be observed as a 50% excess of e^+e^- pairs.

In this Letter we present the analysis of e^+e^- pairs for $m_{ee} < 0.3$ GeV/ c^2 and for $1 < p_T < 5$ GeV/ c in Au + Au and $p + p$ collisions at $\sqrt{s_{NN}} = 200$ GeV recorded during 2004 and 2005, respectively. The PHENIX detector [11] measures electrons in the two central arms, each covering $|\Delta\eta| \leq 0.35$ in pseudorapidity and $\pi/2$ in azimuthal angle. The Au + Au analysis [9,12] uses 8×10^8 minimum bias (Min. Bias) events corresponding to $92.2^{+2.5}_{-3.0}\%$ of the inelastic Au + Au cross section. The beam-beam counters and zero degree calorimeters provide the Min. Bias trigger, as well as the centrality selection [13]. The $p + p$ analysis [14] uses 43 nb^{-1} of data recorded using the Min. Bias trigger and 2.25 pb^{-1} of single electron triggered data. Helium bags in both runs reduced the total conversion material, including the beam pipe, to $\sim 0.4\%$ of a radiation length.

All electrons and positrons with $p_T > 0.2$ GeV/ c are combined into pairs. Pairs from photon conversions in the detector material are removed by a cut on the orientation of the pair in the magnetic field [9]. The combinatorial background is computed by mixing events and is subtracted [9,12]. The S/B ratio is ~ 0.2 (at $m_{ee} = 0.3$ GeV) to ~ 1.5 (at $m_{ee} = 0.1$ GeV/ c^2) for $p_T > 2$ GeV/ c and 0.05 to 0.4

for $1 < p_T < 2$ GeV/c. There are two sources of correlated background: two e^+e^- pairs from a meson decay and correlated hadrons decaying into two e^+e^- pairs, either within the same jet or in back-to-back jets. The magnitude of the correlated background, about 10% of the signal in $p + p$, is determined from the like-sign pair data and subtracted after correcting for acceptance differences between like and unlike-sign pairs [14]. We correct for electron reconstruction efficiency, and in $p + p$ for trigger efficiency, determined as a function of mass and pair p_T using a GEANT-based Monte Carlo simulation [15] of the PHENIX detector.

Figure 1 shows the mass spectra of e^+e^- pairs in $p + p$ and Au + Au collisions for different ranges of pair p_T , comparing to a “cocktail” of hadron decays calculated using a Monte Carlo hadron decay generator based on meson production measured by PHENIX [9]. Detector resolution is included in the cocktail calculation. The open charm contribution, calculated with PYTHIA [16], is also included but is negligible in this kinematic range. The cocktail is normalized to the data for $m_{ee} < 0.03$ GeV/c²; the absolute normalization agrees with the data within a 20% systematic uncertainty [12,14]. The “knee” beginning at $m_{ee} \approx 0.1$ GeV/c² corresponds to the π^0 cutoff, leading to an 80% reduction of background above this point. The $p + p$ data are consistent with the background for $m_{ee} \geq M_{\pi^0}$ at lower p_T , but reveal a small excess over the background at higher p_T . A much greater excess is observed in Au + Au indicating enhanced production of virtual photons.

Internal conversion of direct photons is a possible source of the excess. Little contribution from other sources of e^+e^- pairs is expected in this mass region since $\pi^+\pi^- \rightarrow$

e^+e^- can only contribute for $m_{ee} \geq 2M_\pi$. Although PHENIX has observed a strong enhancement of e^+e^- pairs for $0.15 < m_{ee} < 0.75$ GeV/c² in Au + Au, it peaks at low p_T and decreases rapidly with increasing p_T [9] with a different mass distribution than that observed at high p_T .

Figure 2 shows that the mass spectrum for $m_{ee} < 0.5$ GeV/c² and $p_T > 1$ GeV/c is well described by the cocktail plus internal conversion photons. The flat mass spectrum of the excess above the cocktail at this p_T shows no significant indication of low-mass enhancement [9]. Thus, we treat the excess entirely as internal conversion of photons and deduce the real direct photon yield from e^+e^- pairs using Eq. (1).

We fit a two-component function $f(m_{ee}) = (1 - r)f_c(m_{ee}) + rf_{\text{dir}}(m_{ee})$, to the mass distribution. $f_c(m_{ee})$ is the shape of the cocktail mass distribution (shown in Fig. 1), $f_{\text{dir}}(m_{ee})$ is the expected shape of the direct photon internal conversion, and r is the fit parameter. We assume that the form factor for direct photons is $F(m_{ee}^2) = 1$, as one would expect from a purely pointlike process. For direct photons from parton fragmentation or from hadronic gas, $F(m_{ee}^2)$ may be greater than one. If we arbitrarily set the form factor in $f_{\text{dir}}(m_{ee})$ to be the same as that in $f_\eta(m_{ee})$, r would decrease by $\approx 10\%$.

For each p_T bin, $f(m_{ee})$ is fit to the data for $m_{\text{low}} < m_{ee} < 0.3$ GeV/c² with $m_{\text{low}} = 0.08, 0.1, 0.12$ GeV/c²; r is the only fit parameter. Figure 2 shows $f_{\text{dir}}(m_{ee})$ and $f_c(m_{ee})$ together with a fit result for Au + Au (Min. Bias) data for $1.0 < p_T < 1.5$ GeV/c. For higher p_T bins, χ^2/NDF is near 1.0; fits to centrality separated data also give good χ^2/NDF .

Therefore, we focus on the uncertainties that can cause distortions in the mass distribution: namely, (i) the particle




FIG. 1 (color online). The measured e^+e^- pair invariant mass distributions. The p_T ranges are shown in the legend. The solid curves represent an estimate of hadronic sources; the dashed curves represent the uncertainty in the estimate.




FIG. 2 (color online). Electron pair mass distribution for Au + Au (Min. Bias) events for $1.0 < p_T < 1.5$ GeV/c. The two-component fit is explained in the text. The fit range is $0.12 < m_{ee} < 0.3$ GeV/c². The dashed (black) curve at greater m_{ee} shows $f(m_{ee})$ outside of the fit range.

composition in the hadronic background, (ii) the background (from mixed events and correlated pairs), (iii) the geometric acceptance due to detector active areas, and (iv) the efficiency corrections. These were studied by Monte Carlo simulation. The mass spectrum is distorted within the systematic uncertainties, and the fitting procedure is applied to the distorted spectrum to determine the systematic uncertainties in r . The systematic uncertainty due to the variation of m_{low} is also included. The dominant uncertainty is the particle composition in the hadronic cocktail, namely, the η/π^0 ratio which is $0.48 \pm 0.03(0.08)$ at high p_T for $p + p$ (Au + Au) based on PHENIX measurements [17]. This corresponds to a $\approx 7\%$ ($\approx 17\%$) uncertainty in the $p + p$ (Au + Au) cocktail for $0.1 < m_{ee} < 0.3$ GeV/c². Other sources cause only a few percent uncertainty in the data to cocktail ratio.

Figure 3 shows the fraction r of the direct photon component determined by the two-component fit in (a) $p + p$ and (b) Au + Au (Min. Bias). The curves represent the expectations from a next-to-leading-order perturbative QCD (NLO pQCD) calculation [18]. For $p + p$, the curves show the ratio $d\sigma_\gamma^{NLO}(p_T)/d\sigma_\gamma^{incl}(p_T)$, where $d\sigma_\gamma^{NLO}(p_T)$ is the direct photon cross section from the NLO pQCD calculation and $d\sigma_\gamma^{incl}(p_T)$ is the inclusive photon cross section. For Au + Au, the curves represent $T_{AA}d\sigma_\gamma^{NLO}(p_T)/dN_\gamma^{incl}(p_T)$, where T_{AA} is the Glauber nuclear overlap function and $dN_\gamma^{incl}(p_T)$ is the inclusive photon yield. The three curves correspond, from top to bottom, to the theory scale $\mu = 0.5p_T$, p_T , and $2p_T$, respectively, showing the scale dependence of the theory. While the fraction r is consistent with the NLO pQCD calculation [18] in $p + p$, it is larger than the calculation in Au + Au for $p_T < 3.5$ GeV/c.

The direct photon fraction r in Fig. 3 is converted to the direct photon yield as $dN^{dir}(p_T) = r \times dN^{incl}(p_T)$. The




FIG. 3 (color online). The fraction of the direct photon component as a function of p_T . The error bars and the error band represent the statistical and systematic uncertainties, respectively. The curves are from a NLO pQCD calculation (see text).

inclusive photon yield $dN^{incl}(p_T)$ for each p_T bin is determined from the yield of e^+e^- pairs for $m_{ee} < 0.03$ GeV/c² using Eq. (1). Here we use the fact that in this mass range the process dependent factor S is unity within a few percent for any photon source.

Figure 4 compares the direct photon spectra with previously measured direct photon data from [19,20] and NLO pQCD calculations [18]. The systematic uncertainty of the inclusive photon (14% from the uncertainty in the e^+e^- pair acceptance correction [12]) is added in quadrature with the systematic uncertainties of these data. The $p + p$ data are shown as an invariant cross section using $d\sigma = \sigma_{pp}^{incl} dN$.

In this analysis we have converted the yield of excess e^+e^- pairs to that of real direct photons using Eq. (1), assuming $S = 1$. This implies $\frac{d^2n_{ee}}{dm_{ee}} = \frac{2\alpha}{3\pi} \frac{1}{m_{ee}} dn_\gamma$. Thus the yield of the excess e^+e^- pairs for $0.1 < m_{ee} < 0.3$ GeV/c² before the conversion can be obtained by multiplying the direct photon yield by a factor of $\frac{2\alpha}{3\pi} \log \frac{300}{100} = 1.7 \times 10^{-3}$.

The pQCD calculation is consistent with the $p + p$ data within the theoretical uncertainties for $p_T > 2$ GeV/c. A similarly good agreement is observed for π^0 [21]. The $p + p$ data can be well described by a modified power-law function $[A_{pp}(1 + p_T^2/b)^{-n}]$ as shown by the dashed curve in Fig. 4. The Au + Au data are above the $p + p$ fit curve scaled by T_{AA} for $p_T < 2.5$ GeV/c, indicating that the direct photon yield in the low p_T range increases faster than the binary NN collision scaled $p + p$ cross section.

We fit an exponential plus the T_{AA} -scaled $p + p$ fit function $(Ae^{-p_T/T} + T_{AA} \times A_{pp}(1 + p_T^2/b)^{-n})$ to the Au + Au data. The only free parameters in the fit are A and the inverse slope T of the exponential term. The systematic uncertainties in T are estimated by changing the $p + p$ fit component and the Au + Au data points within the systematic uncertainties. The results of the fits

TABLE I. Summary of the fits. The first and second errors are statistical and systematic, respectively.

Centrality	dN/dy ($p_T > 1$ GeV/c)	T (MeV)	χ^2/DOF
0–20%	$1.50 \pm 0.23 \pm 0.35$	$221 \pm 19 \pm 19$	4.7/4
20–40%	$0.65 \pm 0.08 \pm 0.15$	$217 \pm 18 \pm 16$	5.0/3
Min. Bias	$0.49 \pm 0.05 \pm 0.11$	$233 \pm 14 \pm 19$	3.2/4

are summarized in Table I, where A is converted to dN/dy for $p_T > 1$ GeV/c. For central collisions $T = 221 \pm 19^{\text{stat}} \pm 19^{\text{syst}}$ MeV. Using, instead, a power-law function ($\propto p_T^{-n}$) to fit the $p + p$ spectrum yields $n = 5.40 \pm 0.15$, and $T_{\text{AuAu}} = 240 \pm 21$ MeV. If the direct photons in Au + Au collisions are of thermal origin, the inverse slope T is related to the initial temperature T_{init} of the dense matter. In hydrodynamical models, T_{init} is 1.5 to 3 times T due to the space-time evolution [22]. Several hydrodynamical models can reproduce the central Au + Au data within a factor of 2 [9]. These assume formation of a hot system with initial temperature ranging from $T_{\text{init}} = 300$ MeV at thermalization time $\tau_0 = 0.6$ fm/c to $T_{\text{init}} = 600$ MeV at $\tau_0 = 0.15$ fm/c [22]. As an example, the dotted (red) curve in Fig. 4 shows a thermal photon spectrum in central Au + Au collisions calculated with $T_{\text{init}} = 370$ MeV [7].

In conclusion, we have measured e^+e^- pairs with $m_{ee} < 300$ MeV/c² and $1 < p_T < 5$ GeV/c in $p + p$ and Au +

Au collisions. The $p + p$ data show a small excess over the hadronic background while the Au + Au data show a much larger excess. By treating the excess as internal conversion of direct photons, the direct photon yield is deduced. The yield is consistent with a NLO pQCD calculation in $p + p$. In central Au + Au collisions the shape of the direct photon spectrum above the T_{AA} -scaled $p + p$ spectrum is exponential in p_T , with an inverse slope $T = 221 \pm 19^{\text{stat}} \pm 19^{\text{syst}}$ MeV. Hydrodynamical models with $T_{\text{init}} \sim 300$ –600 MeV at $\tau_0 \sim 0.6$ –0.15 fm/c are in qualitative agreement with the data. Lattice QCD predicts a phase transition from hadronic phase to quark gluon plasma at ~ 170 MeV [1].

We thank the staff of the Collider-Accelerator and Physics Departments at BNL for their vital contributions. We acknowledge support from the Department of Energy and NSF (USA), MEXT and JSPS (Japan), CNPq and FAPESP (Brazil), NSFC (China), MSMT (Czech Republic), IN2P3/CNRS, and CEA (France), BMBF, DAAD, and AvH (Germany), OTKA (Hungary), DAE (India), ISF (Israel), NRF (Korea), MES, RAS, and FAE (Russia), V.R. and KAW (Sweden), U.S. CRDF for the FSU, US-Hungarian NSF-OTKA-MTA, and US-Israel BSF.




FIG. 4 (color online). Invariant cross section ($p + p$) and invariant yield (Au + Au) of direct photons as a function of p_T . The filled points are from this analysis and open points are from [19,20]. The three curves on the $p + p$ data represent NLO pQCD calculations, and the dashed curves show a modified power-law fit to the $p + p$ data, scaled by T_{AA} . The dashed (black) curves are exponential plus the T_{AA} scaled $p + p$ fit. The dotted (red) curve near the 0%–20% centrality data is a theory calculation [7].

*Deceased

†PHENIX Spokesperson:

jacak@skipper.physics.sunysb.edu

- [1] K. Adcox *et al.*, Nucl. Phys. A **757**, 184 (2005).
- [2] K. Adcox *et al.*, Phys. Rev. Lett. **88**, 022301 (2001).
- [3] A. Adare *et al.*, Phys. Rev. Lett. **98**, 172301 (2007).
- [4] K.H. Ackermann *et al.*, Phys. Rev. Lett. **86**, 402 (2001).
- [5] S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 182301 (2003).
- [6] P. Stankus, Annu. Rev. Nucl. Part. Sci. **55**, 517 (2005).
- [7] S. Turbide, R. Rapp, and C. Gale, Phys. Rev. C **69**, 014903 (2004).
- [8] P. Lichard, Phys. Rev. D **51**, 6017 (1995).
- [9] A. Adare *et al.*, arXiv:0911.0244.
- [10] L.G. Landsberg, Phys. Rep. **128**, 301 (1985).
- [11] K. Adcox *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 469 (2003).
- [12] S. Afanasiev *et al.*, arXiv:0706.3034.
- [13] S.S. Adler *et al.*, Phys. Rev. C **69**, 034909 (2004).
- [14] A. Adare *et al.*, Phys. Lett. B **670**, 313 (2009).
- [15] GEANT3.21 CERN Program Library.
- [16] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [17] S.S. Adler *et al.*, Phys. Rev. Lett. **96**, 202301 (2006).
- [18] L.E. Gordon and W. Vogelsang, Phys. Rev. D **48**, 3136 (1993); W. Vogelsang calculated the cross section.
- [19] S.S. Adler *et al.*, Phys. Rev. Lett. **94**, 232301 (2005).
- [20] S.S. Adler *et al.*, Phys. Rev. Lett. **98**, 012002 (2007).
- [21] S.S. Adler *et al.*, Phys. Rev. Lett. **91**, 241803 (2003).
- [22] D. d'Enterria and D. Peressounko, Eur. Phys. J. C **46**, 451 (2006) and references therein.